CrossMark  
click for updates

# Association between cardiometabolic index and kidney stones; a systematic review and meta-analysis

Sina Salem Ahim<sup>1</sup>, Amir Heidari<sup>2</sup>, Leila Ashrafi<sup>3</sup>, Mohammad Rostamzadeh<sup>4</sup>, Negar Jafari<sup>5</sup>, Zahra Eydizadeh<sup>6</sup>, Rasoul Jafari Arismani<sup>7</sup>, Roozbeh Roohinezhad<sup>8</sup>, Abdolmohammad Ranjbar<sup>4</sup>, Mohammad Parsa Mahjoob<sup>9</sup>, Sara Rashki Ghalehno<sup>10\*</sup>

<sup>1</sup>General Medicine, Fasa University of Medical Sciences, Fasa, Iran

<sup>2</sup>Department of Cardiology, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup>Department of Internal Medicine, Clinical Research Development Unit, Hajar Hospital, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>4</sup>Cardiovascular Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>5</sup>Department of Cardiology, School of Medicine, Urmia University of Medical Sciences, Urmia, Iran

<sup>6</sup>Department of Operating Room Nursing, Faculty of Nursing and Midwifery, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

<sup>7</sup>Department of Urologic Surgery, Faculty of Medicine, Arak University of Medical Sciences, Arak, Iran

<sup>8</sup>Department of Urology, Hasheminejad Kidney Center, Iran University of Medical Sciences, Tehran, Iran

<sup>9</sup>Prevention of Cardiovascular Disease Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>10</sup>Department of Cardiology, School of Medicine, Amir al Momenin Hospital, Zabol University of Medical Sciences, Zabol, Iran

## ARTICLE INFO

**Article Type:**  
Meta-analysis

### Article History:

Received: 9 Aug. 2025

Revised: 6 Oct. 2025

Accepted: 4 Nov. 2025

Published online: 18 Dec. 2025

### Keywords:

Cardiometabolic index  
Kidney calculi  
Kidney stone  
Nephrolithiasis  
Renal calculi

## ABSTRACT

**Introduction:** Kidney stone is a common urological disease, and high values of cardiometabolic index (CMI) are primarily linked with metabolic disorders, including obesity, insulin resistance and metabolic syndrome, which are associated with the formation of kidney stones (nephrolithiasis). Accordingly, our study aimed to investigate the relationship between high CMI values and the risk of kidney stones.

**Materials and Methods:** Databases such as Web of Science, Cochrane, Scopus, PubMed, Embase, and Google Scholar Search Engine were conducted for articles published until August 27, 2025. Data were analyzed using STATA 14. Tests with  $P$  values  $< 0.05$  were considered statistically significant.

**Results:** Results revealed that high CMI levels increased the risk of kidney stones (OR=1.39, 95% CI: 1.31, 1.47). As high CMI values in the second one-third (OR=1.38, 95% CI: 1.25, 1.52), third one-third (OR=1.35, 95% CI: 1.22, 1.49), second quartile (OR=1.31, 95% CI: 1.11, 1.56), third quartile (OR=1.45, 95% CI: 1.23, 1.71), and fourth quartile (OR=1.57, 95% CI: 1.33, 1.85) increased the risk of kidney stones. Additionally, high CMI levels increased the risk of kidney stones in men (OR=1.28, 95% CI: 1.16, 1.42), women (OR=1.28, 95% CI: 1.18, 1.39), patients with BMIs 25-30 (OR=1.21, 95% CI: 1.08, 1.35), patients with BMIs  $\geq 30$  (OR=1.25, 95% CI: 1.12, 1.39), patients younger than 60 (OR=1.24, 95% CI: 1.13, 1.37), patients older than 60 (OR=1.36, 95% CI: 1.26, 1.48), and diabetic individuals (OR=1.52, 95% CI: 1.36, 1.70).

**Conclusion:** High CMI levels increased the risk of kidney stone formation by 39%. The risk was similar in men and women; however, the risk of kidney stones increased with the patients' age. Furthermore, obese individuals were at a higher risk compared with those who were overweight.

**Registration:** This study has been compiled in accordance with the PRISMA checklist, and its protocol was registered on the PROSPERO (ID: [CRD420251141496](https://doi.org/10.34172/jrip.2025.38693)) and Research Registry (UIN: [reviewregistry2044](https://doi.org/10.34172/jrip.2025.38693)) websites.

### Implication for health policy/practice/research/medical education:

Our systematic review and meta-analysis found that a high cardiometabolic index (CMI) was a serious risk factor for kidney stones. Studies that grouped CMI levels into quartiles demonstrated that the higher quartiles of CMI were associated with an increased risk of nephrolithiasis compared to the lower quartiles. The patients' gender did not affect the relationship between high CMI values and the risk of kidney stone; hence, male and female patients were equally exposed to the risk of this disease.

**Please cite this paper as:** Salem Ahim S, Heidari A, Ashrafi L, Rostamzadeh M, Jafari N, Eydizadeh Z, Jafari Arismani R, Roohinezhad R, Ranjbar A, Mahjoob MP, Rashki Ghalehno S. Association between cardiometabolic index and kidney stones; a systematic review and meta-analysis. J Renal Inj Prev. 2026; 15(2): e38693. doi: 10.34172/jrip.2025.38693.

## Introduction

Kidney stone is a common urological disease that has affected 10% of the global population (1). In fact, a kidney stone is a chronic metabolic disorder primarily illustrated by the disrupted regulation of salt dissolution and precipitation in urine (2, 3). The incidence and frequency of this disease are increasing around the world, turning kidney stones into a critical public healthcare issue (3, 4). On the other hand, the annual costs of kidney stone diagnosis and treatment surpass billions of dollars, imposing a significant financial burden on individuals and society (5). Besides, kidney stones can cause complications, including acute attacks, urinary tract infections, renal colic, urinary obstruction, and renal failure (6).

The cardiometabolic index (CMI) is an indicator that combines the ratio of triglyceride to high-density lipoprotein-cholesterol (TG/HDL-C) and the ratio of waist-to-height ratio (WHtR) (7). Compared with traditional anthropometric measures, CMI has a closer relationship with metabolic abnormalities as it combines the indicators of abdominal obesity and dyslipidemia (8, 9). Previous studies demonstrated a strong association between high CMI levels and several diseases, including atherosclerosis, ischemic stroke, and hyperuricemia (10-12). On the other hand, high CMI levels are connected with metabolic disorders, such as insulin resistance, metabolic syndrome, and obesity, which are associated with the formation of kidney stones (13, 14).

Since previous studies presented inconsistent results, the present study aimed to examine the relationship between high levels of CMI and the risk of kidney stone using the systematic review and meta-analysis methods; for instance, a study (15) conducted in 2025 reported that, compared with the first quartile, the second quartile did not increase the risk of kidney stones. However, another study (16) in 2025 demonstrated that in the second, third, and fourth quartiles of CMI, the risk of kidney stones was significantly higher than in the first quartile.

## Materials and Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) was used to design the present article (17), and its protocol was registered at the websites PROSPERO (International Prospective Register of Systematic Reviews) and Research Registry.

### Search strategy

The databases such as Web of Science, Cochrane, Scopus, PubMed, Embase, and Google Scholar Search Engine were conducted to search for articles published by August 27, 2025, without any language or publication time restrictions. The Medical Subject Headings (MeSH) and their equivalents were used during the searching process. Operators (AND, OR) were used to combine the keywords. In the end, the process included a manual search. The search strategy in the Embase database was as

follows: 'cardiometabolic index' AND ('kidney calculi' OR 'kidney stone' OR Nephrolithiasis OR 'renal calculi')

### PECO framework

- Population: Articles that aimed to investigate the relationship between high CMI and the risk of kidney stones.
- Exposure: High CMI.
- Comparison: People without kidney stones.
- Outcomes: Risk of kidney stones.

### Inclusion criteria

Articles that aimed to investigate the relationship between high CMI and the risk of kidney stones.

### Exclusion criteria

Case reports, abstracts published in conferences, studies with low qualitative score, duplicate studies, studies that did not have full text, and those that did not provide our required data were excluded.

### Quality assessment

The quality of observational studies was assessed using the Newcastle-Ottawa Scale. This tool assigns a maximum of one star to each question, except for the comparative question. Therefore, a score of zero indicated the lowest quality, and a score of ten showed the highest quality. Then, studies with scores lower than five were considered low-quality (18).

### Data extraction

Two researchers extracted data, including age, stage, number of samples, type of study, country, year, duration of study, and the author's name. Then, the third researcher addressed the discrepancies.

### Statistical analysis

The odds ratio (OR) logarithm was used for data analysis, and the studies were combined. The  $I^2$  index was employed to examine the heterogeneity between studies. A randomized effects model was used to combine the studies. Data analysis was conducted using STATA 14 software. Tests with  $P$  values lower than 0.05 were considered statistically significant.

## Results

Overall, 175 articles were found during the search stage. Then, 83 duplicate studies were identified and removed. The abstracts were reviewed, and 21 studies without accessible full texts were removed. Out of the 71 remaining articles, 45 lacked the required data for analysis and were excluded. Among the 26 articles that entered the next step, 21 studies were removed due to other exclusion criteria, and five articles remained (Figure 1).

As Table 1 shows, five cross-sectional studies with a total of 48625 individuals were investigated, all of which




Figure 1. The PRISMA flow chart of study selection.

Table 1. Summarized information of the studies

Author, year	Country	Type of study	Duration of study	Sample size	Compared with	Stage	OR	Low	Up
Zhong S, 2025 (15)	USA	Cross-sectional	2007–2018	14200	People without kidney stones	Total	1.19	1.06	1.32
						Quartile2	1.16	0.87	1.55
						Quartile3	1.40	1.03	1.91
						Quartile4	1.46	1.08	1.97
Bi J, 2025 (16)	USA	Cross-sectional	2007-2020	18043	People without kidney stones	Total	1.07	1.02	1.12
						Quartile2	1.32	1.01	1.73
						Quartile3	1.44	1.13	1.82
						Quartile4	1.50	1.18	1.92
Liu H, 2024 (19)	USA	Cross-sectional	2007–2018	2714	People without kidney stones	Total	1.17	1.06	1.30
						Quartile2	1.55	1.10	2.18
						Quartile3	1.54	1.09	2.19
						Quartile4	1.88	1.33	2.67
Wang Q, 2024 (20)	USA	Cross-sectional	2011-2018	3059	People without kidney stones	Tertile2	1.39	1.24	1.56
						Tertile3	1.31	1.17	1.47
						Total	1.29	1.16	1.44
Yin G, 2024 (21)	USA	Cross-sectional	2007–2018	10609	People without kidney stones	Tertile2	1.34	1.11	1.62
						Tertile3	1.46	1.20	1.78
						Total	1.29	1.16	1.44

OR: Odds ratio.

were conducted in the United States.

The results revealed that high CMI levels increased the risk of kidney stones (OR=1.39, 95% CI: 1.31, 1.47). As high CMI levels in the second one-third (OR=1.38, 95% CI: 1.25, 1.52), the third one-third (OR=1.35, 95% CI: 1.22, 1.49), the second quartile (OR=1.31, 95% CI: 1.11, 1.56), the third quartile (OR=1.45, 95% CI: 1.23, 1.71), and the

fourth quartile (OR=1.57, 95% CI: 1.33, 1.85) increased the risk of kidney stone formation (Figures 2 and 3).

Moreover, high CMI levels increased the risk of kidney stones in men (OR=1.28, 95% CI: 1.16, 1.42), women (OR=1.28, 95% CI: 1.18, 1.39), patients with body mass indexes of 25 to 30 kg/m<sup>2</sup> (OR=1.21, 95% CI: 1.08, 1.35), and in patients with body mass indexes of ≥ 30 kg/m<sup>2</sup>




Figure 2. Forest plot showing the association between CMI and kidney stones.




Figure 3. Forest plot showing the association between CMI and kidney stones by stage.

(OR=1.25, 95% CI: 1.12, 1.39) (Figures 4 to 7).


Additionally, high CMI levels increased the risk of patients younger than 60 years (OR=1.24, 95% CI: 1.13, 1.37), older than 60 years (OR=1.36, 95% CI: 1.26, 1.48), and diabetic individuals (OR=1.52, 95% CI: 1.36, 1.70) (Figures 8 to 10).

**Discussion**


Based on the findings of the current meta-analysis, high CMI levels increased the risk of kidney stones (39%). As high CMI levels in the second one-third (38%), third one-third (35%), second quartile (31%), third quartile (45%), and fourth quartile (57%) increased the risk of kidney stone formation. Besides, high CMI levels increased the risk of kidney stones in men (28%), women (28%), patients with BMIs 25-30 (21%), patients with BMIs ≥ 30 (25), patients younger than 60 (24%), patients older than

60 (36%), and diabetic patients (52%).


In a study by Bi et al high CMI levels were associated with increased risk of kidney stones (OR= 1.07, 95% CI: 1.02, 1.12) (16). Furthermore, Liu et al demonstrated that high CMI levels were positively correlated with increased risk of kidney stones in diabetic patients (OR= 1.17, 95% CI: 1.06, 1.30) (19). According to the results of Wang et al the odds ratios for the occurrence of kidney stones in the second and the third one-third were (OR= 1.39, 95% CI: 1.24, 1.56) and (OR= 1.31, 95% CI: 1.17, 1.47), respectively (20). In another cross-sectional study by Yin et al on 10609 participants, researchers reported that high CMI levels led to the formation of kidney stones (OR= 1.29, 95% CI: 1.16, 1.44) (21). Zhong et al demonstrated a statistically significant relationship between high CMI levels and the increased frequency of kidney stone formation (OR= 1.19, 95% CI: 1.06, 1.32) (15). The mentioned studies




**Figure 4.** Forest plot showing the association between CMI and kidney stones in males.




**Figure 7.** Forest plot showing the relationship between CMI and kidney stones in individuals with body mass index ≥30 kg/m<sup>2</sup>.




**Figure 5.** Forest plot showing the association between CMI and kidney stones in females.



**Figure 8.** Forest plot showing the relationship between CMI and kidney stones in individuals with mean age <60 years.



**Figure 6.** Forest plot showing the relationship between CMI and kidney stones in individuals with body mass index (25-30 kg/m<sup>2</sup>).



**Figure 9.** Forest plot showing the relationship between CMI and kidney stones in individuals with mean age ≥60 years.



- G, Lange D. Mechanisms of the intestinal and urinary microbiome in kidney stone disease. *Nat Rev Uro.* 2022 Dec;19:695-707. doi: 10.1038/s41585-022-00647-5
2. Gao H, Lin J, Xiong F, Yu Z, Pan S, Huang Y. Urinary microbial and metabolomic profiles in kidney stone disease. *Front Cell Infect Microbiol.* 2022;12:953392. doi: 10.3389/fcimb.2022.953392
  3. Khan S, Pearle M, Robertson W, Gambaro G, Canales B, Doizi S, et al. Kidney stones. *Nat Rev Dis Primers.* 2016;2:16008. doi: 10.1038/nrdp.2016.8
  4. Singh P, Harris PC, Sas DJ, Lieske JC. The genetics of kidney stone disease and nephrocalcinosis. *Nat Rev Nephrol.* 2022;18:224-40. doi: 10.1038/s41581-021-00513-4.
  5. Anderegg M, Olinger E, Bargagli M, Geraghty R, Taylor L, Nater A, et al. Prevalence and characteristics of genetic disease in adult kidney stone formers. *Nephrol Dial Transplant.* 2024 Sep;39:1426-41. doi: 10.1093/ndt/gfae074
  6. Ganesan C, Holmes M, Liu S, Montez-Rath M, Conti S, Chang T, et al. Kidney stone events after kidney transplant in the United States. *Clin J Am Soc Nephrol.* 2023;18:777-84. doi: 10.2215/CJN.000000000000176
  7. Zhou X, Tao X, Zhang L, Yang Q, Li Z, Dai L, et al. Association between cardiometabolic index and depression: National Health and Nutrition Examination Survey (NHANES) 2011–2014. *J Affect Disord.* 2024;351:939-47. doi: 10.1016/j.jad.2024.02.024
  8. Siener R. Nutrition and Kidney Stone Disease Nutrients. 2021 Jun 13:1917. doi: 10.3390/nu13061917
  9. Zha F, Cao C, Hong M, Hou H, Zhang Q, Tang B, et al. The nonlinear correlation between the cardiometabolic index and the risk of diabetes: A retrospective Japanese cohort study. *Front Endocrinol (Lausanne).* 2023;14:1120277. doi: 10.3389/fendo.2023.1120277
  10. Cai X, Hu J, Wen W, Wang J, Wang M, Liu S, et al. Associations of the Cardiometabolic Index with the risk of Cardiovascular Disease in patients with hypertension and obstructive sleep apnea: results of a Longitudinal Cohort Study. *Oxid Med Cell Longev.* 2022;2022:4914791. doi: 10.1155/2022/4914791.
  11. Li FE, Luo Y, Zhang FL, Zhang P, Liu D, Ta S, et al. Association between cardiometabolic index and stroke: A population-based cross-sectional study. *Current neurovascular research.* 2021;18:324-32. doi: 10.2174/1567202618666211013123557.
  12. Zuo Y, Gao Z, Yin Y, Yang X, Feng P. Association between the cardiometabolic index and hyperuricemia in an asymptomatic population with normal body mass index. *Int J Gen Med.* 2021;14:8603-10. doi: 10.2147/IJGM.S340595
  13. Song J, Li Y, Zhu J, Liang J, Xue S, Zhu Z. Non-linear associations of cardiometabolic index with insulin resistance, impaired fasting glucose, and type 2 diabetes among US adults: a cross-sectional study. *Front Endocrinol (Lausanne).* 2024;15:1341828. doi: 10.3389/fendo.2024.1341828
  14. Duan S, Yang D, Xia H, Ren Z, Chen J, Yao S. Cardiometabolic index: A new predictor for metabolic associated fatty liver disease in Chinese adults. *Front Endocrinol (Lausanne).* 2022;13:1004855. doi: 10.3389/fendo.2022.1004855
  15. Zhong S, Li G, Tian C, Jiang M, Chen D, Wang H, et al. Associations between cardiometabolic index with kidney stones: evidence from NHANES 2007–2018. *Front Endocrinol (Lausanne).* 2025;16:1485477. doi: 10.3389/fendo.2025.1485477
  16. Bi J, Du J, Yan X, Chen R. Diabetes mediates an inverted L-shaped association between cardiometabolic index and kidney stones: a cross-sectional study from NHANES 2007–2020. *Front Endocrinol (Lausanne).* 2025;16:1535724. doi: 10.3389/fendo.2025.1535724
  17. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev J.* 2015;4:1e9. doi: 10.1186/2046-4053-4-1.
  18. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010;25:603e5.
  19. Liu H, Guo M, Zhou Y, Jin M, Hao H, Yuan Y, et al. Association between cardiometabolic index and kidney stones in US diabetic patients: NHANES 2007–2018. *World J Urol.* 2024;42:653. doi: 10.1007/s00345-024-05364-9
  20. Wang Q, Wang Z, Tang Z, Liu C, Pan Y, Zhong S. Association between cardiometabolic index and kidney stone from NHANES: a population-based study. *Front Endocrinol (Lausanne).* 2024;15:1408781. doi: 10.3389/fendo.2024.1408781
  21. Yin G, Zhang W, Zhang J, Sheng T, Chen B. Diabetes mediates the relationship between cardiometabolic index and kidney stones: a cross-sectional study. *Sci Rep.* 2024;14:31075. doi: 10.1038/s41598-024-82075-6
  22. Kong J, Tao W, Sun Y, Xu Y, Li H, Li J. The relationship between cardiometabolic index and diabetic kidney disease in people with diabetes. *Front Endocrinol (Lausanne).* 2025;15:1376813. doi: 10.3389/fendo.2024.1376813
  23. Xu Q, Lin J, Liao L, Hu J, Lu J. Association between cardiometabolic index and albuminuria: evidence from NHANES 2017–2020. *PloS One.* 2025;20:e0318736. doi: 10.1371/journal.pone.0318736
  24. Guo Q, Wang Y, Liu Y, Wang Y, Deng L, Liao L, et al. Association between the cardiometabolic index and chronic kidney disease: a cross-sectional study. *Int Urol Nephrol.* 2024;56:1733-41. doi: 10.1007/s11255-023-03888-4
  25. Zhu Z, Ren K, Duan X, Hu X, Lv Y, Wang D, et al. Cardiometabolic index is associated with impaired renal function: Nonlinear threshold effects and dose-response relationships in a national cohort (NHANES 1999–2018). *Medicine.* 2025;104:e43653. doi: 10.1097/MD.00000000000043653
  26. Guo T, Zhou Y, Yang G, Sheng L, Chai X. Association between cardiometabolic index and hypertension among US adults from NHANES 1999–2020. *Sci Rep.* 2025;15:4007. doi: 10.1038/s41598-025-87029-0